


Example of ARCALYST® (rilonacept) Coverage Decision Logic for Recurrent Pericarditis*



RP = recurrent pericarditis; PA = prior authorization; DMARD = Disease modifying antirheumatic drug; TB = tuberculosis

*This an example of common payer coverage decision making; however, each payer may differ in the determination of coverage for ARCALYST. Please refer to specific payer policies on ARCALYST access requirements.

INDICATION

ARCALYST is indicated for the treatment of recurrent pericarditis (RP) and reduction in risk of recurrence in adults and pediatric patients 12 years and older.

IMPORTANT SAFETY INFORMATION

Warnings and Precautions

- Interleukin-1 (IL-1) blockade may interfere with the immune response to infections. Treatment with another medication that works through inhibition of IL-1 or inhibition of tumor necrosis factor (TNF) is not recommended as this may increase the risk of serious infection. Serious, life-threatening infections have been reported in patients taking ARCALYST. Do not initiate treatment with ARCALYST in patients with an active or chronic infection.

Please see additional Important Safety Information on next page.



IMPORTANT SAFETY INFORMATION (Cont.)

Warnings and Precautions (Cont.)

- Discontinue ARCALYST if a patient develops a serious infection.
- It is possible that taking drugs such as ARCALYST that block IL-1 may increase the risk of tuberculosis (TB) or other atypical or opportunistic infections.
- Although the impact of ARCALYST on infections and the development of malignancies is not known, treatment with immunosuppressants, including ARCALYST, may result in an increase in the risk of malignancies.
- Hypersensitivity reactions associated with ARCALYST occurred in clinical trials. Discontinue ARCALYST and initiate appropriate therapy if a hypersensitivity reaction occurs.
- Increases in non-fasting lipid profile parameters occurred in patients treated with ARCALYST in clinical trials. Patients should be monitored for changes in their lipid profiles.
- Since no data are available, avoid administration of live vaccines while patients are receiving ARCALYST. ARCALYST may interfere with normal immune response to new antigens, so vaccines may not be effective in patients receiving ARCALYST. It is recommended that, prior to initiation of therapy with ARCALYST, patients receive all recommended vaccinations, as appropriate.

Adverse Reactions

- The most common adverse reactions ($\geq 10\%$) include injection-site reactions and upper respiratory tract infections.

Drug Interactions

- In patients being treated with CYP450 substrates with narrow therapeutic indices, therapeutic monitoring of the effect or drug concentration should be performed, and the individual dose of the medicinal product may need to be adjusted.

For more information about ARCALYST, see [Full Prescribing Information](#).



ARCALYST is a registered trademark of Regeneron Pharmaceuticals, Inc.
Kiniksa OneConnect is a trademark of Kiniksa Pharmaceuticals.
©2024 Kiniksa Pharmaceuticals. All Rights Reserved. 06/24 ARC-US-00172-24

